

1	SHORT COMMUNICATION	63
2		64
3		65
4	Energy metabolism and cellular homeostasis trade-offs provide	66
5	the basis for a new type of sensitivity to ocean acidification in a	67
6	marine polychaete at a high CO₂ vent: adenylate and phosphagen	68
7	energy pools versus carbonic anhydrase	69
8	Lucy M. Turner^{1,2,*}, Elena Ricevuto³, Alexia Massa Gallucci³, Maria-Cristina Gambi³ and Piero Calosi^{1,4}	70
9		71
10		72
11		73
12		74
13		75
14		76
15		77
16	ABSTRACT	78
17	Species distributions and ecology can often be explained by their	79
18	physiological sensitivity to environmental conditions. Whilst we have	80
19	a relatively good understanding of how these are shaped by	81
20	temperature, for other emerging drivers, such as P _{CO₂} we know	82
21	relatively little. The marine polychaete <i>Sabella spallanzanii</i> increases	83
22	its metabolic rate when exposed to high P _{CO₂} conditions and remains	84
23	absent from the CO ₂ vent of Ischia. To understand new possible	85
24	pathways of sensitivity to CO ₂ in marine ectotherms, we examined the	86
25	metabolic plasticity of <i>S. spallanzanii</i> exposed <i>in situ</i> to elevated	87
26	P _{CO₂} by measuring fundamental metabolite and carbonic anhydrase	88
27	concentrations . We show that whilst this species can survive elevated	89
28	P _{CO₂} conditions in the short term, and exhibits an increase in energy	90
29	metabolism, this is accompanied by a significant decrease in	91
30	carbonic anhydrase concentration . These homeostatic changes are	92
31	unlikely to be sustainable in the longer term, indicating <i>S. spallanzanii</i>	93
32	may struggle with future high P _{CO₂} conditions.	94
33		95
34	KEY WORDS: Individual approach, Ocean acidification, Climate	96
35	change, Homeostatic capacity, Trade-offs, Mediterranean Sea	97
36		98
37	INTRODUCTION	99
38	An understanding of a species' ecophysiological limits helps us	100
39	identify the reasons underpinning species and population processes	101
40	over different spatial scales. Whilst we have a relatively good	102
41	understanding of how species distributions are shaped by	103
42	temperature (see Bozinovic et al., 2011), for other emerging	104
43	environmental drivers, such as increasing seawater partial pressure	105
44	of CO₂ (P_{CO₂}) , which is causing the acidification of the oceans, we	106
45	know relatively little (see Calosi et al., 2013; Kroeker et al., 2011;	107
46	Maas et al., 2012). Shallow water, high CO ₂ vents have been used as	108
47	analogue to investigate the potential ecological and evolutionary	109
48	implications of ocean acidification. In particular, at the CO ₂ vent of	110
49	Ischia the polychaete fauna have been characterised in relation to the	111
50	venting activity (e.g. Kroeker et al., 2011). Based on their	112
51	distribution patterns, species found inside and outside these	113
52	naturally acidified areas can be considered to be either 'tolerant'	114
53	(abundant both inside and outside the low pH/high P _{CO₂} areas) or	115
54		116
55		117
56	¹ Marine Biology and Ecology Research Centre, Plymouth University, Drake Circus,	118
57	Plymouth PL4 8AA, UK. ² Department of Biological and Environmental Sciences,	119
58	University of Gothenburg, Box 461, Gothenburg 405 30, Sweden. ³ Stazione	120
59	Zoologica Anton Dohrn, Villa Dohrn, Ischia, Napoli, Italy. ⁴ Université du Québec à	121
	Rimouski, Rimouski, Quebec, Canada G5L 3A1.	122
60	*Author for correspondence (lucy.turner@gu.se)	123
61		124
62	Received 1 December 2014; Accepted 11 May 2015	1