

## RESEARCH ARTICLE

## Benthic protists: the under-charted majority

Dominik Forster<sup>1</sup>, Micah Dunthorn<sup>1</sup>, Frédéric Mahé<sup>1,†</sup>, John R. Dolan<sup>2</sup>, Stéphane Audic<sup>3</sup>, David Bass<sup>4,5</sup>, Lucie Bittner<sup>1,3,6</sup>, Christophe Boutte<sup>3</sup>, Richard Christen<sup>7,8</sup>, Jean-Michel Claverie<sup>9</sup>, Johan Decelle<sup>3</sup>, Bente Edvardsen<sup>10</sup>, Elianne Egge<sup>10</sup>, Wenche Eikrem<sup>10</sup>, Angélique Gobet<sup>3,11</sup>, Wiebe H.C.F. Kooistra<sup>12</sup>, Ramiro Logares<sup>13</sup>, Ramon Massana<sup>13</sup>, Marina Montresor<sup>12</sup>, Fabrice Not<sup>3</sup>, Hiroyuki Ogata<sup>2,14</sup>, Jan Pawlowski<sup>15</sup>, Massimo C. Pernice<sup>13</sup>, Sarah Romac<sup>3</sup>, Kamran Shalchian-Tabrizi<sup>10</sup>, Nathalie Simon<sup>3</sup>, Thomas A. Richards<sup>16</sup>, Sébastien Santini<sup>9</sup>, Diana Sarno<sup>12</sup>, Raffaele Siano<sup>17</sup>, Daniel Vaultot<sup>3</sup>, Patrick Wincker<sup>18</sup>, Adriana Zingone<sup>12</sup>, Colomán de Vargas<sup>3</sup> and Thorsten Stoeck<sup>1,\*</sup>

<sup>1</sup>Department of Ecology, University of Kaiserslautern, Erwin-Schrödinger Str. 14, D-67663 Kaiserslautern, Germany, <sup>2</sup>Sorbonne Universités, UPMC Univ. Paris 06, CNRS, UMR CNRS UMR 7093 and Laboratoire d'Océanographie de Villefranche-sur-Mer, Université Paris 06, F-06230 Villefranche-sur-Mer, France, <sup>3</sup>Sorbonne Universités, UPMC Univ. Paris 06, CNRS, UMR 7144, Station Biologique, Place Georges Teissier, 29680 Roscoff, France, <sup>4</sup>Department of Life Sciences, the Natural History Museum London, Cromwell Road, London SW7 5BD, UK, <sup>5</sup>Centre for Environment, Fisheries and Aquaculture Science (Cefas), Barrack Road, the Nothe, Weymouth, Dorset DT4 8UB, UK, <sup>6</sup>Sorbonne Universités, UPMC, CNRS, Institut de Biologie Paris-Seine (IBPS), Evolution Paris Seine, F-75005 Paris, France, <sup>7</sup>CNRS, UMR 7138 & Université de Nice-Sophia Antipolis, F-06103 Nice cedex 2, France, <sup>8</sup>Université de Nice-Sophia Antipolis & CNRS, UMR 7138 F-06103 Nice cedex 2, France, <sup>9</sup>CNRS, Aix-Marseille Université, IGS UMR7256, F-13288 Marseille, France, <sup>10</sup>Department of BioSciences, University of Oslo, Blindern, 0316 N-Oslo, Norway, <sup>11</sup>CNRS, UMR 8227 & UPMC Université Paris 06, Station Biologique de Roscoff, F-29682 Roscoff, France, <sup>12</sup>Stazione Zoologica Anton Dohrn, Villa Comunale 1, I-80121, Naples, Italy, <sup>13</sup>Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37–49, ES-08003, Barcelona, Catalonia, Spain, <sup>14</sup>Institute for Chemical Research, Kyoto University, Uji, 611-0011, Japan, <sup>15</sup>Department of Genetics and Evolution, University of Geneva, 4, Boulevard d'Yvoy, CH-1211 Geneva, Switzerland, <sup>16</sup>Biosciences, University of Exeter, Exeter, EX4 4QD, UK, <sup>17</sup>Ifremer, Centre de Brest DYNECO/Pelagos Technopôle Brest Iroise, BP 7029280 Plouzané, France and <sup>18</sup>CEA, Genoscope, 2 rue Gaston Crémieux, F-91000 Evry, France

\*Corresponding author: Department of Ecology, University of Kaiserslautern, Erwin-Schroedinger-Straße 14, D-67633 Kaiserslautern, Germany.

Tel: +49-631-2052502; Fax: +49-631-2052496; E-mail: [stoeck@rhrk.uni-kl.de](mailto:stoeck@rhrk.uni-kl.de)

**One sentence summary:** Sampling RNA in near-shore marine environments along the European coast shows that benthic protist communities are extremely diverse with many species currently unknown.

Editor: Julie Olson

†Frédéric Mahé, <http://orcid.org/0000-0002-2808-0984>

Received: 11 March 2016; Accepted: 30 May 2016

© FEMS 2016. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)