



## Review

## The Whittard Canyon – A case study of submarine canyon processes



T. Amaro<sup>a,†</sup>, V.A.I. Huvenne<sup>b</sup>, A.L. Allcock<sup>c</sup>, T. Aslam<sup>d,e</sup>, J.S. Davies<sup>f</sup>, R. Danovaro<sup>g,h</sup>, H.C. De Stigter<sup>i</sup>, G.C.A. Duineveld<sup>i</sup>, C. Gambi<sup>g</sup>, A.J. Gooday<sup>b</sup>, L.M. Gunton<sup>b</sup>, R. Hall<sup>d</sup>, K.L. Howell<sup>f,j</sup>, J. Ingels<sup>j</sup>, K. Kiriakoulakis<sup>k</sup>, C.E. Kershaw<sup>k</sup>, M.S.S. Lavaley<sup>i</sup>, K. Robert<sup>b</sup>, H. Stewart<sup>l</sup>, D. Van Rooij<sup>m</sup>, M. White<sup>c</sup>, A.M. Wilson<sup>c</sup>

<sup>a</sup> Hellenic Center for Marine Research (HCMR), 710 03 Heraklion, Crete, Greece

<sup>b</sup> National Oceanography Centre, University of Southampton, Waterfront Campus, Southampton SO14 3ZH, UK

<sup>c</sup> Ryan Institute and School of Natural Sciences, National University of Ireland, Galway, University Road, Galway, Ireland

<sup>d</sup> Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich, UK

<sup>e</sup> Centre for Environment Fisheries and Aquaculture Sciences (Cefas), Lowestoft, UK

<sup>f</sup> Marine Biology & Ecology Research Centre, Marine Institute, Plymouth University, Plymouth PL4 8AA, UK

<sup>g</sup> Dep Life and Environmental Sciences, Polytechnic University of Marche, Ancona, Italy

<sup>h</sup> Stazione Zoologica Anton Dohrn, Villa Comunale I, Napoli, Italy

<sup>i</sup> NIOZ Royal Netherlands Institute for Sea Research, Department of Ocean Systems Sciences, and Utrecht University, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

<sup>j</sup> Plymouth Marine Laboratory, Prospect Place, West Hoe, PL1 3DH Plymouth, UK

<sup>k</sup> School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool L3 3AF, UK

<sup>l</sup> British Geological Survey, The Lyell Centre, Research Avenue South, Edinburgh EH9 3LA, UK

<sup>m</sup> Renard Centre of Marine Geology (RCMG), Department of Geology and Soil Science, Ghent University, Krijgslaan 281 S8, B-9000 Gent, Belgium

## article info

## Article history:

Received 5 December 2015

Received in revised form 13 June 2016

Accepted 13 June 2016

Available online 18 June 2016

## abstract

Submarine canyons are large geomorphological features that incise continental shelves and slopes around the world. They are often suggested to be biodiversity and biomass hotspots, although there is no consensus about this in the literature. Nevertheless, many canyons do host diverse faunal communities but owing to our lack of understanding of the processes shaping and driving this diversity, appropriate management strategies have yet to be developed. Here, we integrate all the current knowledge of one single system, the Whittard Canyon (Celtic Margin, NE Atlantic), including the latest research on its geology, sedimentology, geomorphology, oceanography, ecology, and biodiversity in order to address this issue. The Whittard Canyon is an active system in terms of sediment transport. The net suspended sediment transport is mainly up-canyon causing sedimentary overflow in some upper canyon areas. Occasionally sediment gravity flow events do occur, some possibly the result of anthropogenic activity. However, the role of these intermittent gravity flows in transferring labile organic matter to the deeper regions of the canyon appears to be limited. More likely, any labile organic matter flushed downslope in this way becomes strongly diluted with bulk material and is therefore of little food value for benthic fauna. Instead, the fresh organic matter found in the Whittard Channel mainly arrives through vertical deposition and lateral transport of phytoplankton blooms that occur in the area during spring and summer. The response of the Whittard Canyon fauna to these processes is different in different groups. Foraminiferal abundances are higher in the upper parts of the canyon and on the slope than in the lower canyon. Meiofaunal abundances in the upper and middle part of the canyon are higher than on adjacent slopes, but lower in the deepest part. Mega- and macrofauna abundances are higher in the canyon compared with the adjacent slope and are higher in the eastern than the western branch. These faunal patterns reflect the fact that the Whittard Canyon encompasses considerable environmental heterogeneity, related to a combination of organic matter trapping, current regimes (due to focused internal tides) and different substrates. We conclude that coordinated observations of processes driving faunal patterns are needed at a fine scale in order to understand the functioning of communities in this and other submarine canyons.

© 2016 Elsevier Ltd. All rights reserved.

<sup>†</sup> Corresponding author.