

Roles of Retinoic Acid in Cardiomyocytes Specification during *Ciona* Embryogenesis

Director of Studies: Enrico D'Aniello

Department Biology and Evolution of Marine Organisms

Seat: Naples

Abstract

Vitamin A (retinol) and its derivatives play several essential roles during embryogenesis and throughout adult life in Metazoa, acting both at the genetic and epigenetic levels. The earliest requirement of all-trans retinoic acid (atRA) in the development of vertebrate deuterostomes occurs in the posteriorization of the body axis, which affects the patterning of several organs, including the spinal cord, forelimbs, eye, reproductive tracts and heart. In vertebrates, RA signalling is necessary for proper vertebrate heart development as either decrease or increase of RA embryonic levels can result in congenital heart malformations. To date, retinoic acid signalling pathway has been shown to exist in ascidians, and to have general developmental functions conserved with vertebrates. However, specific and detailed studies of the role of RA signalling during heart development are still missing in tunicates. The ascidian *Ciona* has emerged as a significant model system for the study of chordate development including the heart. **Therefore, given the high level of conservation with vertebrates of the key genes involved in cardiac cell specification and differentiation, the goal of this PhD project will be to determine the requirements and the role of RA signalling for proper heart development during *Ciona* embryogenesis.** The use of a non-canonical model system such as *Ciona* has the potential to dramatically improve our understanding of inappropriate RA signalling that will ultimately provide new insight to shed light also to human congenital heart defects and disease.